1/9/1
DIALOG(R)File 347:JAPIO
(c) 2002 JPO & JAPIO. All rts. reserv. 04381388

PEPTIDE OR ITS SALT

PUB. NO.: 06-025288 [JP 6025288 A] PUBLISHED: February 01, 1994 (19940201)

INVENTOR(s): TANIHARA MASAO

FUJIWARA CHIE

APPLICANT(s): KURARAY CO LTD [000108] (A Japanese Company or Corporation), JP (Japan)

APPL. NO.: 04-207501 [JP 92207501] FILED: July 10, 1992 (19920710)

INTL CLASS: [5] C07K-007/08; C07K-007/06; C07K-007/10; A61K-037/02;

A61K-037/02; A61K-037/02; A61K-037/02; A61K-037/02;

C07K-099/00

JAPIO CLASS: 14.1 (ORGANIC CHEMISTRY -- Organic Compounds); 14.4 (ORGANIC CHEMISTRY -- Medicine)

JOURNAL: Section: C, Section No. 1196, Vol. 18, No. 236, Pg. 91, May 06, 1994 (19940506)

ABSTRACT

PURPOSE: To obtain a peptide, useful for treating fracture, wound, multiple sclerosis, rheumatoid arthritis, osteoarthritis, osteoporosis, periodontosis, etc., and capable of effectively suppressing the rejection after organ transplantation.

CONSTITUTION: This peptide or its salt has the following general formula: H- X.up arrow.1-Ala-X.up arrow.2-Pro-Cys-Cys-Val-X.up arrow.3-Gln-X.up arrow.4-Leu-Glu-X.up arrow.5-Y (X.up arrow.2 denotes an amino acid residue selected from the group consisting of Ser and Ala; X.up arrow.3 denotes an amino acid residue selected from the group consisting of Ser and Pro; X.up arrow.4 denotes an amino acid residue selected from the group consisting of Ala and Asp; X.up arrow.1 and X.up arrow.5 respectively denote single bonds or peptide fragments composed of 1-5 amino acid residues selected from the group consisting of Gly, Ala, Val, Arg, Asn, Ser, Phe, Pro, Leu, Glu, Asp, Lys, Thr, His, Tyr, Nle and Ile; Y denotes hydroxyl group or amino group) and transforming growth factor (TGF)-.beta.-like activity.